

## FOLIAR FEEDING PROGRAMME FOR CABBAGE, CAULIFOLWER, BROCCOLI, BRUSSEL SPROUTS



Growth Stage	Product	Rate / ha	Comments
<b>Trans-planter Barrel</b>	MaxiBoost Multifeed Magnisal or MagMax	1 ℓ 2 kg 500 g	Apply in sufficient water at planting time
<b>2 – 3 Weeks after planting to sufficient foliage</b>	MaxiBoost Multifeed Magnisal or MagMax	1 ℓ 3 kg 1 kg or 2 ℓ	Apply in 300 – 500 ℓ water / ha. Reduce product rate and water volumes accordingly for row application.
<b>Every 7 – 10 days throughout growing season</b>	Supafeed Magnisal or MagMax Spraybor CalciMax	4 kg 1 kg or 2ℓ 3 – 4 ℓ 300 – 500 g 1,5 – 3 ℓ	Apply in 300 – 500 ℓ water / ha. Reduce product rate and water volume accordingly for row application.
<b>After fruit starts to size up</b>	Polyfeed Magnisal or MagMax Spraybor CalciMax	3 kg 1 kg or 2 ℓ 3 – 5 ℓ 300 – 400 g 1,5 – 3 ℓ	Make 3 – 4 applications in 300 – 500 ℓ water / ha, 7 – 10 days apart.
<b>When heads size up until harvest</b>	CalciMax Spraybor	5 ℓ 300 – 500 g	2 – 3 Applications in 300 – 500 ℓ water / ha, 7 – 10 days apart. Adequate calcium supplied by 20 – 30 ℓ CalciMax / ha / season. This will help prevent split head and maintain firmness.